

09/369883

Att #14

WEST Search History

DATE: Thursday, May 22, 2003

| <u>Set Name</u> side by side | <u>Query</u> | <u>Hit Count</u> | <u>Set Name</u> result set |
|--|-------------------------|------------------|-------------------------------|
| <i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ</i> | | | |
| L4 | l1 with L3 | 236 | L4 |
| L3 | brij\$ or tween\$ | 40938 | L3 |
| L2 | liposom\$ or lipofect\$ | 41594 | L2 |
| L1 | liposom\$ or lipofect\$ | 41594 | L1 |

END OF SEARCH HISTORY

WEST**Print Selection**[Help](#)[Clear](#)[Cancel](#)[Print](#)[Print First Page](#)

| Select? | Document ID | Section(s) | Page(s) | # Pages to print | Database |
|-------------------------------------|---------------|------------|---------|------------------|--------------------------|
| <input checked="" type="checkbox"/> | 6451603 | all | all | 25 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 6559123 | all | all | * 86 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 6471956 | all | all | * 153 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 6432917 | all | all | 47 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 5942237 | all | all | 9 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 5030442 | all | all | 27 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | WO009639121A1 | all | all | N/A | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 4330534 | all | all | 4 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 4937078 | all | all | 7 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 5492924 | all | all | 10 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 4861597 | all | all | 14 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | 5053217 | all | all | 9 | USPT,PGPB,JPAB,EPAB,DWPI |
| <input checked="" type="checkbox"/> | US5820880A | all | all | 14 | USPT,PGPB,JPAB,EPAB,DWPI |

Note: Print requests for more than 49 pages are denoted by '*' and are in red.

Building**Room****Printer**

cm1



11e14



gbldptr

[Main Menu](#)[Logout](#)

09/369883
Act #14

=> s brij? or tween?
L1 33443 BRIJ? OR TWEEN?

=> s liposom?
L2 122173 LIPOSOM?

=> s l1 and l2

L3 435 L1 AND L2

=> dup rem l3
PROCESSING COMPLETED FOR L3
L4 255 DUP REM L3 (180 DUPLICATES REMOVED)

=> s l4 and py<1997
2 FILES SEARCHED...
4 FILES SEARCHED...
L5 132 L4 AND PY<1997

=> d l5 ibib abs 1-132

L5 ANSWER 1 OF 132 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
ACCESSION NUMBER: 1997:65259 BIOSIS
DOCUMENT NUMBER: PREV199799364462
TITLE: New cationic lipid formulations for gene transfer.
AUTHOR(S): Liu, Feng; Yang, Jingping; Huang, Leaf; Liu, Dexi (1)
CORPORATE SOURCE: (1) Dep. Pharm. Sci., Sch. Pharm., Univ. Pittsburgh,
Pittsburgh, PA 15261 USA
SOURCE: Pharmaceutical Research (New York), (1996) Vol. 13, No. 12,
pp. 1856-1860.
ISSN: 0724-8741.
DOCUMENT TYPE: Article
LANGUAGE: English
AB Purpose: To develop appropriate dosage forms of DNA for gene delivery.

Methods: 3-beta(N-(N,N'-dimethylaminoethane) carbamoyl) cholesterol (DC-Chol) was mixed either with ***Tween*** 80 alone, or with additional lipid components including castor oil and phosphatidylcholine (PC) or dioleoylphosphatidylethanolamine (DOPE) to make different lipid formulations. The particle size and the physical stability of the formulations upon mixing with plasmid DNA containing the luciferase cDNA

were examined using laser light scattering measurement. The transfection activity of the DNA/lipid complexes was tested in presence or absence of serum using a cell culture system. Results: We demonstrated that many favorable properties as a gene carrier could be achieved by formulating DNA into new dosage forms using ***Tween*** 80 as the major emulsifier. Compared to the cationic ***liposomes***, these new formulations transfected different cell lines with an equivalent or higher efficiency. Not only are they resistant to serum, but also form stable DNA complexes which could be stored for longer periods of time without losing transfection activity. Conclusions: Cationic lipids formulated into different lipid formulations using ***Tween*** 80 as a surfactant appeared to have more favorable physical and biological activities than traditional cationic ***liposomes*** as a carrier for gene delivery.

L5 ANSWER 2 OF 132 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
ACCESSION NUMBER: 1997:31549 BIOSIS
DOCUMENT NUMBER: PREV199799337952
TITLE: Lyophilized preliposomal formulation of the non-cross-resistant anthracycline annamycin: Effect of surfactant on ***liposome*** formation, stability and size.
AUTHOR(S): Zou, Yiyu (1); Priebe, Waldemar; Perez-Soler, Roman
CORPORATE SOURCE: (1) Sect. Exp. Ther., Dep. Thoracic/Head Neck Med. Oncol.,
Univ. Texas M.D. Anderson Cancer Cent., 1515 Holcombe Blvd., Houston, TX 77030 USA
SOURCE: Cancer Chemotherapy and Pharmacology, (1996) Vol. 39, No. 1-2, pp. 103-108.
ISSN: 0344-5704.
DOCUMENT TYPE: Article

LANGUAGE: English

AB We report a method of preparing a submicron and stable ***liposome***

formulation of the non-cross-resistant anthracycline annamycin. The lipids were dimyristoylphosphatidyl choline (DMPC) and dimyristoylphosphatidyl glycerol (DMPG) at a 7:3 molar ratio and the optimal lipid: drug ratio was 50:1 (w/w). The selected formulation was a preliposome lyophilized powder

that contained the phospholipids, annamycin, and ***Tween*** 20. The ***liposome*** suspension was obtained on the day of use by adding normal saline at 37 degree C (1 ml/mg annamycin) and hand shaking for 1 min. The presence of ***Tween*** 20 was essential in shortening the reconstitution step (from gt 2 h to 1 min), avoiding the early formation of free drug crystals, and reducing the median particle size by tenfold (from 1.5 mu-m to 0.15 mu-m) without destroying the ***liposome*** vesicles. At room temperature, the preliposome powder was chemically stable for gt 3 months, and the ***liposome*** suspension was chemically and physically stable for gt 24 h. The in vitro cytotoxicity of the formulation was equivalent to that of the same lipid composition prepared by the standard evaporation method. The results of the study indicate that small amounts of surfactant may be used to enhance the reconstitution step and reduce the size of ***liposome*** suspensions obtained from lyophilized preliposome powders. The formulation described is being used for ongoing clinical trials with ***liposomal*** annamycin.

L5 ANSWER 3 OF 132 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
ACCESSION NUMBER: 1996:465538 BIOSIS
DOCUMENT NUMBER: PREV199699187894
TITLE: Intrathecal chemotherapy with 1,3-Bis(2-chloroethyl)-1-nitrosourea encapsulated into hybrid ***liposomes*** for meningeal gliomatosis: An experimental study.
AUTHOR(S): Kitamura, Isao (1); Kochi, Masato; Matsumoto, Yoko; Ueoka,

Ryuichi; Kuratsu, Jun-Ichi; Ushio, Yukioka
CORPORATE SOURCE: (1) Dep. Neurosurg., Kumamoto Univ. Med. Sch., 1-1-1 Honjo,
Kumamoto 860 Japan
SOURCE: Cancer Research, (1996) Vol. 56, No. 17, pp. 3986-3992.
ISSN: 0008-5472.

DOCUMENT TYPE: Article
LANGUAGE: English

AB 1,3-bis(2-chloroethyl)-1-nitrosourea (BCNU), one of the chloroethyl nitrosoureas, is effective against malignant glioma. To develop its use in intrathecal chemotherapy, we encapsulated BCNU in hybrid ***liposomes*** composed of dimyristoylphosphatidylcholine and micellar surfactants (***Tween*** 20) and dissolved it in artificial cerebrospinal fluid (lipo-BCNU). We then studied the toxicity of hybrid ***liposomes*** and cellular proliferation inhibition of lipo-BCNU in vitro. We found that 3 mM hybrid ***liposomes*** did not affect the viability of human endothelial cells and that lipo-BCNU inhibited the proliferation of human glioma cell lines U-105MG, U-251MG, and U-373MG, and rat glioma cell lines

C6 and 9L in a concentration-dependent fashion. Wistar rats that were administered lipoBCNU intracisternally showed no weight loss, neurological symptoms, or histological changes of the brain and spinal cord. A Wistar rat model of meningeal gliomatosis was established by intracisternal inoculation of 0.1 ml cell suspension containing 1 times 10⁻⁶ or 5 times 10⁻⁶ viable C6 glioma cells. Two days after inoculation, lipo-BCNU (BCNU,

2.5 mg/kg) was administered intracisternally. When 1 times 10⁻⁶ glioma cells were inoculated (experiments 1 and 2), the median survival times were 24.5 and 26 days in the control groups and 32 and 45 days in the lipo-BCNU-treated groups, respectively. When 5 times 10⁻⁶ glioma cells were inoculated (experiments 3-6), the median survival times were 17-29.5

days in the control groups and 23-44 days in the treated groups, respectively. Significantly prolonged survival was obtained in three of six experimental groups. After the administration of 1 ml lipo-BCNU (BCNU, 4.67 mM) or 1 ml BCNU solubilized with 5% dextrose/water (BCNU, 4.67 mM)

into the cisterna magna of dogs, the cisterna magna cerebrospinal fluid